Neuroscience Letters, 176 (1994) 282–285
Elsevier Ireland Ltd.

AUTHOR INDEX

Acosta, G.B. and Rubio, M.C., $GABA_A$ receptors mediate the changes produced by stress on GABA function and locomotor activity, 29
Alberghina, M., Infarinato, S., Anfuso, C.D., Lupo, G. and Sipione, S., Lipid hydroperoxides induce changes in palmitate uptake across the rat blood–retina and blood–brain barrier, 247
Albuquerque, E.X., see Alkondon, M., 152
Alkondon, M. and Albuquerque, E.X., Presence of α-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat olfactory bulb neurons, 152
Altagracia, M., Monroy-Noyola, A., Osorio-Rico, L., Kravzov, J., Alvarado-Calvillo, R., Manjarrez-Marmolejo, J. and Ríos, C., Dapsone attenuates kainic acid-induced seizures in rats, 52
Alvarado-Calvillo, R., see Altagracia, M., 52
Ambrosio, S., see Unzeta, M., 235
Anderson, J.J., Kuo, S. and Chase, T.N., Endogenous excitatory amino acids tonically stimulate striatal acetylcholine release through NMDA but not AMPA receptors, 264
Anfuso, C.D., see Alberghina, M., 247
Araki, N., see Ohta, K., 165
Araki, T., Kato, H., Liu, X.-H., Itoyama, Y., Kogure, K. and Kato, K., Delayed damage of striatal interneurons after cerebral ischemia in the gerbil, 17
Arckens, L., see Zhang, F., 137
Arias, J.L., see Lopez, L., 197
Arimura, A., see Légrádi, G., 193
Asou, H., see Yazaki, T., 13
Austin, D.R., see Gold, B.G., 123

Baas, D., Fressinaud, C., Ittel, M.E., Reeber, A., Dalençon, D., Puymirat, J. and Sarlième, L.L., Expression of thyroid hormone receptor isoforms in rat oligodendrocyte cultures. Effect of 3,5,3′-triiodo-L-thyronine, 47
Babey, A.M., Palmour, R.M. and Young, S.N., Caffeine and propranolol block the increase in rat pineal melatonin production produced by stimulation of adenosine receptors, 93
Bagetta, G., Finazzi-Agrò, A., Palma, E. and Nisticò, G., Intracerebral injection of human immunodeficiency virus type 1 coat glycoprotein GP120 does not produce neurodegeneration in rats, 97
Baldessarini, R.J., Campbell, A., Ben-Jonathan, N., Ellingboe, J., Zong, R. and Neumeyer, J.L., Effects of aporphine isomers on rat prolactin, 269
Baron, S., see Unzeta, M., 235
Bauer, M., see McLachlan, E.M., 147
Beaulieu, C., see Morin, F., 85
Ben-Jonathan, N., see Baldessarini, R.J., 269
Berti, F., see Rossoni, G., 185
Bigiani, A. and Roper, S.D., Reduction of electrical coupling between *Necturus* taste receptor cells, a possible role in acid taste, 212
Birbaumer, N., see Lutzenberger, W., 115
Bohn, M.C., see Phillipson, O.T., 67
Bothmer, J., Markerink, M. and Jolles, J., Phosphoinositide kinase activities in synaptosomes prepared from brains of patients with Alzheimer's disease and controls, 169
Braña, M., see Lopez, L., 197
Brookes, A., see Clair, D.S., 45
Brösamle, C., see Skatchkov, S., 251
Brown, G.M., see Kapur, S., 21
Burgos, P., see Lopez, L., 197

Calancie, B., see Klose, K.J., 277
Campbell, A., see Baldessarini, R.J., 269
Cass, W.A. and Gerhardt, G.A., Direct in vivo evidence that D2 dopamine receptors can modulate dopamine uptake, 259
Chaouloff, F., Influence of physical exercise on $5\text{-}HT_{1A}$ receptor- and anxiety-related behaviours, 226
Chapman, V. and Dickenson, A.H., Enhanced responses of rat dorsal horn neurones after UV irradiation of the hindpaw; roles of the NMDA receptor, 41
Chase, T.N., see Anderson, J.J., 264
Clair, D.S., Norrman, J., Perry, R., Yates, C., Wilcock, G. and Brookes, A., Apolipoprotein E ε4 allele frequency in patients with Lewy body dementia, Alzheimer's disease and age-matched controls, 45
Clementi, F., see Rossoni, G., 185
Constanti, A., see Harris, N.C., 221
Cornish, J.L. and den Buuse, M.v., Pressor responses to electrical and chemical stimulation of the rat brain A10 dopaminergic system, 142
Cummings, J.A., Nicola, S.M. and Malenka, R.C., Induction in the rat hippocampas of long-term potentiation (LTP) and long-term depression (LTD) in the presence of a nitric oxide synthase inhibitor, 110

Dalençon, D., see Baas, D., 47
Dalmaz, Y., see Poncet, L., 1
de Blas, A., see Krantis, A., 32
De Koninck, Y., Salter, M.W. and Henry, J.L., Substance P released endogenously by high-intensity sensory stimulation potentiates purinergic inhibition of nociceptive dorsal horn neurons induced by peripheral vibration, 128
Denoroy, L., see Poncet, L., 1
Dickenson, A.H., see Chapman, V., 41
Domann, R., Westerhoff, C.H.A. and Witte, O.W., Inhibitory mechanisms terminating paroxysmal depolarization shifts in hippocampal neurons of rats, 71

Ellingboe, J., see Baldessarini, R.J., 269

Finazzi-Agrò, A., see Bagetta, G., 97
Fressinaud, C., see Baas, D., 47
Frotscher, M., see Plaschke, M., 243
Fukunari, A., Kato, A., Sakai, Y., Yoshimoto, T., Ishiura, S., Suzuki, K. and Nakajima, T., Colocalization of prolyl endopeptidase and amyloid β-peptide in brains of senescence-accelerated mouse, 201
Fukushima, N., Ueda, H., Hayashi, C., Katayama, T., Miyamae, T. and Misu, Y., Species and age-dependent differences of functional coupling between opioid δ-receptor and G-proteins and possible involvement of protein kinase C in striatal membranes, 55
Fukuuchi, Y., see Kobari, M., 157
Fukuuchi, Y., see Ohta, K., 165
Furukawa, S., see Nakajima, M., 161

Galazka, P., see Ravindran, J., 209
Gendrot, G., see Servière, J., 80
Gentleman, S.M., see Griffin, W.S.T., 133
Gerhardt, G.A., see Cass, W.A., 259
Gold, B.G., Austin, D.R. and Storm-Dickerson, T., Multiple signals

Neuroscience Letters, 176 (1994) 282–285
Elsevier Ireland Ltd.

AUTHOR INDEX

Acosta, G.B. and Rubio, M.C., $GABA_A$ receptors mediate the changes produced by stress on GABA function and locomotor activity, 29
Alberghina, M., Infarinato, S., Anfuso, C.D., Lupo, G. and Sipione, S., Lipid hydroperoxides induce changes in palmitate uptake across the rat blood–retina and blood–brain barrier, 247
Albuquerque, E.X., see Alkondon, M., 152
Alkondon, M. and Albuquerque, E.X., Presence of α-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat olfactory bulb neurons, 152
Altagracia, M., Monroy-Noyola, A., Osorio-Rico, L., Kravzov, J., Alvarado-Calvillo, R., Manjarrez-Marmolejo, J. and Ríos, C., Dapsone attenuates kainic acid-induced seizures in rats, 52
Alvarado-Calvillo, R., see Altagracia, M., 52
Ambrosio, S., see Unzeta, M., 235
Anderson, J.J., Kuo, S. and Chase, T.N., Endogenous excitatory amino acids tonically stimulate striatal acetylcholine release through NMDA but not AMPA receptors, 264
Anfuso, C.D., see Alberghina, M., 247
Araki, N., see Ohta, K., 165
Araki, T., Kato, H., Liu, X.-H., Itoyama, Y., Kogure, K. and Kato, K., Delayed damage of striatal interneurons after cerebral ischemia in the gerbil, 17
Arckens, L., see Zhang, F., 137
Arias, J.L., see Lopez, L., 197
Arimura, A., see Légrádi, G., 193
Asou, H., see Yazaki, T., 13
Austin, D.R., see Gold, B.G., 123

Baas, D., Fressinaud, C., Ittel, M.E., Reeber, A., Dalençon, D., Puymirat, J. and Sarlième, L.L., Expression of thyroid hormone receptor isoforms in rat oligodendrocyte cultures. Effect of 3,5,3′-triiodo-L-thyronine, 47
Babey, A.M., Palmour, R.M. and Young, S.N., Caffeine and propranolol block the increase in rat pineal melatonin production produced by stimulation of adenosine receptors, 93
Bagetta, G., Finazzi-Agrò, A., Palma, E. and Nisticò, G., Intracerebral injection of human immunodeficiency virus type 1 coat glycoprotein GP120 does not produce neurodegeneration in rats, 97
Baldessarini, R.J., Campbell, A., Ben-Jonathan, N., Ellingboe, J., Zong, R. and Neumeyer, J.L., Effects of aporphine isomers on rat prolactin, 269
Baron, S., see Unzeta, M., 235
Bauer, M., see McLachlan, E.M., 147
Beaulieu, C., see Morin, F., 85
Ben-Jonathan, N., see Baldessarini, R.J., 269
Berti, F., see Rossoni, G., 185
Bigiani, A. and Roper, S.D., Reduction of electrical coupling between *Necturus* taste receptor cells, a possible role in acid taste, 212
Birbaumer, N., see Lutzenberger, W., 115
Bohn, M.C., see Phillipson, O.T., 67
Bothmer, J., Markerink, M. and Jolles, J., Phosphoinositide kinase activities in synaptosomes prepared from brains of patients with Alzheimer's disease and controls, 169
Braña, M., see Lopez, L., 197
Brookes, A., see Clair, D.S., 45
Brösamle, C., see Skatchkov, S., 251
Brown, G.M., see Kapur, S., 21
Burgos, P., see Lopez, L., 197

Calancie, B., see Klose, K.J., 277
Campbell, A., see Baldessarini, R.J., 269
Cass, W.A. and Gerhardt, G.A., Direct in vivo evidence that D2 dopamine receptors can modulate dopamine uptake, 259
Chaouloff, F., Influence of physical exercise on $5\text{-}HT_{1A}$ receptor- and anxiety-related behaviours, 226
Chapman, V. and Dickenson, A.H., Enhanced responses of rat dorsal horn neurones after UV irradiation of the hindpaw; roles of the NMDA receptor, 41
Chase, T.N., see Anderson, J.J., 264
Clair, D.S., Norrman, J., Perry, R., Yates, C., Wilcock, G. and Brookes, A., Apolipoprotein E ε4 allele frequency in patients with Lewy body dementia, Alzheimer's disease and age-matched controls, 45
Clementi, F., see Rossoni, G., 185
Constanti, A., see Harris, N.C., 221
Cornish, J.L. and den Buuse, M.v., Pressor responses to electrical and chemical stimulation of the rat brain A10 dopaminergic system, 142
Cummings, J.A., Nicola, S.M. and Malenka, R.C., Induction in the rat hippocampas of long-term potentiation (LTP) and long-term depression (LTD) in the presence of a nitric oxide synthase inhibitor, 110

Dalençon, D., see Baas, D., 47
Dalmaz, Y., see Poncet, L., 1
de Blas, A., see Krantis, A., 32
De Koninck, Y., Salter, M.W. and Henry, J.L., Substance P released endogenously by high-intensity sensory stimulation potentiates purinergic inhibition of nociceptive dorsal horn neurons induced by peripheral vibration, 128
Denoroy, L., see Poncet, L., 1
Dickenson, A.H., see Chapman, V., 41
Domann, R., Westerhoff, C.H.A. and Witte, O.W., Inhibitory mechanisms terminating paroxysmal depolarization shifts in hippocampal neurons of rats, 71

Ellingboe, J., see Baldessarini, R.J., 269

Finazzi-Agrò, A., see Bagetta, G., 97
Fressinaud, C., see Baas, D., 47
Frotscher, M., see Plaschke, M., 243
Fukunari, A., Kato, A., Sakai, Y., Yoshimoto, T., Ishiura, S., Suzuki, K. and Nakajima, T., Colocalization of prolyl endopeptidase and amyloid β-peptide in brains of senescence-accelerated mouse, 201
Fukushima, N., Ueda, H., Hayashi, C., Katayama, T., Miyamae, T. and Misu, Y., Species and age-dependent differences of functional coupling between opioid δ-receptor and G-proteins and possible involvement of protein kinase C in striatal membranes, 55
Fukuuchi, Y., see Kobari, M., 157
Fukuuchi, Y., see Ohta, K., 165
Furukawa, S., see Nakajima, M., 161

Galazka, P., see Ravindran, J., 209
Gendrot, G., see Servière, J., 80
Gentleman, S.M., see Griffin, W.S.T., 133
Gerhardt, G.A., see Cass, W.A., 259
Gold, B.G., Austin, D.R. and Storm-Dickerson, T., Multiple signals

underlie the axotomy-induced up-regulation of c-JUN in adult sensory neurons, 123
Gold, P.E., see Walker, D.L., 255
Gold, R., see Zettl, U.K., 75
Graham, D.I., see Griffin, W.S.T., 133
Griffin, W.S.T., Sheng, J.G., Gentleman, S.M., Graham, D.I., Mrak, R.E. and Roberts, G.W., Microglial interleukin-1α expression in human head injury: correlations with neuronal and neuritic β-amyloid precursor protein expression, 133
Guerra, G., see Martínez, A., 239

Halleux, P., see Zhang, F., 137
Hamada, J., see Ohta, K., 165
Harris, N.C., Libri, V. and Constanti, A., Selective blockade of the hyperpolarization-activated cationic current (I_h) in guinea pig substantia nigra pars compacta neurones by a novel bradycardic agent, Zeneca ZM 227189, 221
Hartung, H.-P., see Zettl, U.K., 75
Hayashi, C., see Fukushima, N., 55
Hayashi, K., see Nakajima, M., 161
Hayashi, Y., see Nakajima, M., 161
Henry, J.L., see De Koninck, Y., 128
Hirano, A., see Kawanami, T., 178
Houle, S., see Kapur, S., 21
Hutcheon, B., see Puil, E., 63

Ijaz, S., see Ravindran, J., 209
Infarinato, S., see Alberghina, M., 247
Ishaqzay, R., see Ravindran, J., 209
Ishiura, S., see Fukunari, A., 201
Itoyama, Y., see Araki, T., 17
Ittel, M.E., see Baas, D., 47
Izumi, Y. and Zorumski, C.F., Developmental changes in the effects of metabotropic glutamate receptor antagonists on CA1 long-term potentiation in rat hippocampal slices, 89

Jolles, J., see Bothmer, J., 169
Jouvet, M., see Poncet, L., 1

Kaba, H., see Li, C.-S., 5
Kachi, T., see Suzuki, T., 217
Kapur, S., Meyer, J., Wilson, A.A., Houle, S. and Brown, G.M., Activation of specific cortical regions by apomorphine: an $[^{15}O]H_2O$ PET study in humans, 21
Kashiwagi, K., see Nakajima, M., 161
Kasper, E.M., see Plaschke, M., 243
Katayama, T., see Fukushima, N., 55
Kato, A., see Fukunari, A., 201
Kato, H., see Araki, T., 17
Kato, K., see Araki, T., 17
Kato, T., see Kawanami, T., 178
Kawanami, T., Kato, T., Llena, J.F., Hirano, A. and Sasaki, H., Altered synaptophysin-immunoreactive pattern in human olivary hypertrophy, 178
Kawashima, T., see Nakajima, M., 161
Keast, J.R., see McLachlan, E.M., 147
Kim, H., see Takahashi, L.K., 272
Klose, K.J. and Calancie, B., Has GM1 ganglioside been shown to be effective in the restoration of neurologic function in persons with chronic spinal cord injury? A critique of an article by Judith B. Walker and Michelle Harris, 277
Kobari, M., Fukuuchi, Y., Tomita, M., Tanahashi, N., Konno, S. and Takeda, H., Dilatation of cerebral microvessels mediated by endothelin ET_B receptor and nitric oxide in cats, 157
Kogure, K., see Araki, T., 17
Komatsumoto, S., see Ohta, K., 165
Konno, S., see Kobari, M., 157
Krantis, A., Nichols, K., de Blas, A. and Staines, W., Demonstration of benzodiazepine receptors in submucosal neurons of the gastrointestinal tract, 32
Kravzov, J., see Altagracia, M., 52
Kuffler, D.P., see Skatchkov, S., 251
Kuo, S., see Anderson, J.J., 264

La Maestra, L., see Rossoni, G., 185
Layer, P.G., see Weikert, T., 9
Légrádi, G., Shioda, S. and Arimura, A., Pituitary adenylate cyclase-activating polypeptide-like immunoreactivity in autonomic regulatory areas of the rat medulla oblongata, 193
Lerchl, A., Increased oxidation of pineal serotonin as a possible explanation for reduced melatonin synthesis in the aging Djungarian hamster (*Phodopus sungorus*), 25
LeSauter, J., see Servière, J., 80
Li, C.-S., Kaba, H. and Seto, K., Effective induction of pregnancy block by electrical stimulation of the mouse accessory olfactory bulb coincident with prolactin surges, 5
Li, P., see Zhang, Y.M., 231
Libri, V., see Harris, N.C., 221
Liu, L., see Ravindran, J., 209
Liu, X.-H., see Araki, T., 17
Llena, J.F., see Kawanami, T., 178
Lopez, L., Braña, M., Burgos, P. and Arias, J.L., Structural dimorphism in the mammillary bodies of the rat, 197
Lovick, T.A., see Zhang, Y.M., 231
Lupo, G., see Alberghina, M., 247
Lutzenberger, W., Pulvermüller, F. and Birbaumer, N., Words and pseudowords elicit distinct patterns of 30-Hz EEG responses in humans, 115

Mahy, N., see Unzeta, M., 235
Mailleux, P., see Zhang, F., 137
Malenka, R.C., see Cummings, J.A., 110
Manjarrez-Marmolejo, J., see Altagracia, M., 52
Markerink, M., see Bothmer, J., 169
Martínez, A., Muñoz-Clares, R.A., Guerra, G., Morán, J. and Pasantes-Morales, H., Sulfhydryl groups essential for the volume-sensitive release of taurine from astrocytes, 239
Matsuyama, S., see Murakami, N., 181
McLachlan, E.M., Keast, J.R. and Bauer, M., SP- AND CGRP-immunoreactive axons differ in their ability to reinnervate the skin of the rat tail, 147
Meyer, J., see Kapur, S., 21
Misu, Y., see Fukushima, N., 55
Miura, M., see Yazaki, T., 13
Miura, M., see Takayama, K., 59
Miyamae, T., see Fukushima, N., 55
Miyashita, H., see Ravindran, J., 209
Moneta, M.E., see Strecker, R.E., 173
Monroy-Noyola, A., see Altagracia, M., 52
Morán, J., see Martínez, A., 239
Morin, F. and Beaulieu, C., Equivalent cell density in three areas of neonatal rat cerebral cortex, 85
Mrak, R.E., see Griffin, W.S.T., 133
Mullikin-Kilpatrick, D. and Treistman, S.N., Time course of nerve

growth factor modulation of ethanol inhibition of Ca^{2+} currents in PC12 cells, 101
Muñoz-Clares, R.A., see Martínez, A., 239
Murakami, N., Sakai, N., Nei, K., Matsuyama, S., Saito, N. and Tanaka, C., Potassium and calcium channel involvement in induction of long-lasting synaptic enhancement by calyculin A, a protein phosphatase inhibitor, in rat hippocampal CA1 region, 181

Nakajima, M., Kashiwagi, K., Ohta, J., Furukawa, S., Hayashi, K., Kawashima, T. and Hayashi, Y., Nerve growth factor and epidermal growth factor rescue PC12 cells from programmed cell death induced by etoposide: distinct modes of protection against cell death by growth factors and a protein-synthesis inhibitor, 161
Nakajima, T., see Fukunari, A., 201
Nakamura, H., see Yamada, Y., 189
Naumann, T., see Plaschke, M., 243
Nei, K., see Murakami, N., 181
Neumeyer, J.L., see Baldessarini, R.J., 269
Nichols, K., see Krantis, A., 32
Nicola, S.M., see Cummings, J.A., 110
Nisticò, G., see Bagetta, G., 97
Norrman, J., see Clair, D.S., 45

Oh, Y., Sashihara, S. and Waxman, S.G., In situ hybridization localization of the Na^+ channel $\beta 1$ subunit mRNA in rat CNS neurons, 119
Ohta, J., see Nakajima, M., 161
Ohta, K., Araki, N., Shibata, M., Hamada, J., Komatsumoto, S., Shimazu, K. and Fukuuchi, Y., A novel in vivo assay system for consecutive measurement of brain nitric oxide production combined with the microdialysis technique, 165
Okada, Y., see Yamada, Y., 189
Orban, G.A., see Zhang, F., 137
Orkand, R.K., see Skatchkov, S., 251
Osorio-Rico, L., see Altagracia, M., 52

Palma, E., see Bagetta, G., 97
Palmour, R.M., see Babey, A.M., 93
Pasantes-Morales, H., see Martínez, A., 239
Péquignot, J.-M., see Poncet, L., 1
Perez, V., see Unzeta, M., 235
Perry, R., see Clair, D.S., 45
Phillipson, O.T. and Bohn, M.C., C1–3 adrenergic medullary neurones project to the paraventricular thalamic nucleus in the rat, 67
Plaschke, M., Kasper, E.M., Naumann, T. and Frotscher, M., Survival and transmitter expression of rat cholinergic medial septal neurons despite removal of hippocampus in the early postnatal period, 243
Poncet, L., Denoroy, L., Dalmaz, Y., Péquignot, J.-M. and Jouvet, M., Chronic hypoxia affects peripheral and central vasoactive intestinal peptide-like immunoreactivity in the rat, 1
Puil, E., Hutcheon, B. and Reiner, P.B., Isoflurane inhibits calcium currents in neocortical neurons, 63
Pulvermüller, F., see Lutzenberger, W., 115
Puymirat, J., see Baas, D., 47

Ravindran, J., Shuaib, A., Ijaz, S., Galazka, P., Waqar, T., Ishaqzay, R., Miyashita, H. and Liu, L., High extracellular GABA levels in hippocampus – as a mechanism of neuronal protection in cerebral ischemia in adrenalectomized gerbils, 209
Reeber, A., see Baas, D., 47
Reiner, P.B., see Puil, E., 63
Ríos, C., see Altagracia, M., 52
Roberts, G.W., see Griffin, W.S.T., 133
Roper, S.D., see Bigiani, A., 212
Rossoni, G., Berti, F., La Maestra, L. and Clementi, F., ω-Conotoxin GVIA binds to and blocks rat neuromuscular junction, 185
Rubio, M.C., see Acosta, G.B., 29

Saito, N., see Murakami, N., 181
Sakai, N., see Murakami, N., 181
Sakai, Y., see Fukunari, A., 201
Salter, M.W., see De Koninck, Y., 128
Sarlièvе, L.L., see Baas, D., 47
Sasaki, H., see Kawanami, T., 178
Sashihara, S., see Oh, Y., 119
Schousboe, A., see Westergaard, N., 105
Servière, J., Gendrot, G., LeSauter, J. and Silver, R., Host resets phase of grafted SCN: influence of implant site, tissue specificity and pineal secretion, 80
Seto, K., see Li, C.-S., 5
Sheng, J.G., see Griffin, W.S.T., 133
Shibata, M., see Ohta, K., 165
Shieh, G.-J. and Walters, D.E., Altered neurochemical and behavioral development of 10-day-old rats perinatally exposed to the κ opioid agonist U-50,488H, 37
Shimazu, K., see Ohta, K., 165
Shioda, S., see Légrádi, G., 193
Shuaib, A., see Ravindran, J., 209
Silver, R., see Servière, J., 80
Simon, R.P., see Tanaka, K., 205
Sipione, S., see Alberghina, M., 247
Skatchkov, S., Brõsamle, C., Vyklický, L., Kuffler, D.P. and Orkand, R.K., NMDA receptors on adult frog spinal motoneurons in culture, 251
Sonnewald, U., see Westergaard, N., 105
Staines, W., see Krantis, A., 32
Storm-Dickerson, T., see Gold, B.G., 123
Strecker, R.E. and Moneta, M.E., Electrical stimulation of the kindled hippocampus briefly increases extracellular dopamine in the nucleus accumbens, 173
Suzuki, K., see Fukunari, A., 201
Suzuki, T. and Kachi, T., Differences between adrenaline and noradrenaline cells in cellular association with supporting cells in the adrenal medulla of the pig: an immunohistochemical study, 217
Suzuki, T., see Takayama, K., 59

Takahashi, L.K. and Kim, H., Intracranial action of corticosterone facilitates the development of behavioral inhibition in the adrenalectomized preweanling rat, 272
Takayama, K., Suzuki, T. and Miura, M., The comparison of effects of various anesthetics on expression of Fos protein in the rat brain, 59
Takeda, H., see Kobari, M., 157
Tanahashi, N., see Kobari, M., 157
Tanaka, C., see Murakami, N., 181
Tanaka, K. and Simon, R.P., The pattern of neuronal injury following seizures induced by intranigral kainic acid, 205
Tomita, M., see Kobari, M., 157
Toya, S., see Yazaki, T., 13
Toyka, K.V., see Zettl, U.K., 75
Treistman, S.N., see Mullikin-Kilpatrick, D., 101

Ueda, H., see Fukushima, N., 55
Unzeta, M., Baron, S., Perez, V., Ambrosio, S. and Mahy, N., Sex-related effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydro-pyridine treatment may be related to differences in monoamine oxidase B, 235

Uyemura, K., see Yazaki, T., 13

Van Brée, L., see Zhang, F., 137
Van den Buuse, M., see Cornish, J.L., 142
Vanderhaeghen, J.-J., see Zhang, F., 137
Vandesande, F., see Zhang, F., 137
Vanduffel, W., see Zhang, F., 137
Vyklický, L., see Skatchkov, S., 251

Walker, D.L. and Gold, P.E., Intra-amygdala kinase inhibitors disrupt retention of a learned avoidance response in rats, 255
Walters, D.E., see Shieh, G.-J., 37
Waqar, T., see Ravindran, J., 209
Waxman, S.G., see Oh, Y., 119
Weikert, T. and Layer, P.G., The carbohydrate epitope HNK-1 is present on all inactive, but not on all active forms of chicken butyrylcholinesterase, 9
Westergaard, N., Sonnewald, U. and Schousboe, A., Release of α-ketoglutarate, malate and succinate from cultured astrocytes: possible role in amino acid neurotransmitter homeostasis, 105
Westerhoff, C.H.A., see Domann, R., 71
Wilcock, G., see Clair, D.S., 45
Wilson, A.A., see Kapur, S., 21
Witte, O.W., see Domann, R., 71

Yamada, Y., Nakamura, H. and Okada, Y., Propentofylline enhances the formation of long-term potentiation in guinea pig hippocampal slices, 189
Yates, C., see Clair, D.S., 45
Yazaki, T., Miura, M., Asou, H., Toya, S. and Uyemura, K., Peripheral myelin P0 protein mediates neurite outgrowth of cortical neurons in vitro and axonal regeneration in vivo, 13
Yoshimoto, T., see Fukunari, A., 201
Young, S.N., see Babey, A.M., 93

Zettl, U.K., Gold, R., Hartung, H.-P. and Toyka, K.V., Apoptotic cell death of T-lymphocytes in experimental autoimmune neuritis of the Lewis rat, 75
Zhang, F., Halleux, P., Arckens, L., Vanduffel, W., Van Brée, L., Mailleux, P., Vandesande, F., Orban, G.A. and Vanderhaeghen, J.-J., Distribution of immediate early gene zif-268, c-fos, c-jun and jun-D mRNAs in the adult cat with special references to brain region related to vision, 137
Zhang, Y.M., Li, P. and Lovick, T.A., Role of the nucleus raphe obscurus in the inhibition of rostral ventrolateral medullary neurones induced by stimulation in the ventrolateral periaqueductal grey matter of the rabbit, 231
Zong, R., see Baldessarini, R.J., 269
Zorumski, C.F., see Izumi, Y., 89